

Form Inequalities for Symmetric Contraction Semigroups

Markus Haase

Abstract. Consider — for the generator $-A$ of a symmetric contraction semigroup over some measure space X , $1 \leq p < \infty$, q the dual exponent and given measurable functions $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ — the statement:

$$\operatorname{Re} \sum_{j=1}^m \int_X A F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \geq 0$$

for all $\mathbb{C}^d$ -valued measurable functions $\mathbf{f}$ on X such that $F_j(\mathbf{f}) \in \operatorname{dom}(A_p)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all j .

It is shown that this statement is valid in general if it is valid for X being a two-point Bernoulli $(\frac{1}{2}, \frac{1}{2})$ -space and A being of a special form. As a consequence we obtain a new proof for the optimal angle of L^p -analyticity for such semigroups, which is essentially the same as in the well-known sub-Markovian case.

The proof of the main theorem is a combination of well-known reduction techniques and some representation results about operators on $C(K)$ -spaces. One focus of the paper lies on presenting these auxiliary techniques and results in great detail.

Mathematics Subject Classification (2010). 47A60, 47D06, 47D07, 47A07.

Keywords. symmetric contraction semigroup, diffusion semigroup, sector of analyticity, Stone model, integral bilinear forms, tensor products.

Second Version, 29 August, 2015.

1. Introduction

In the recent preprint [2], A. Carbonaro and O. Dragičević consider symmetric contraction semigroups $(S_t)_{t \geq 0}$ over some measure space $X = (X, \Sigma, \mu)$ and prove so-called spectral multiplier results (= functional calculus estimates) for A_p , where $-A_p$ is the generator of $(S_t)_{t \geq 0}$ on $L^p(X)$, $1 \leq p < \infty$.

Part of this work was supported by the Marsden Fund Council from Government funding, administered by the Royal Society of New Zealand.

Their proof consists of three major steps. In the first one, the authors show how to generate functional calculus estimates for the operator $A = A_p$ from *form inequalities* of the type

$$\sum_{j=1}^m \operatorname{Re} \int_X [AF_j(f_1, \dots, f_d)] \cdot G_j(f_1, \dots, f_d) \, d\mu \geq 0, \quad (1.1)$$

where F_j and G_j are measurable functions $\mathbb{C}^d \rightarrow \mathbb{C}$ with certain properties and $(f_1, \dots, f_d)$ varies over a suitable subset of measurable functions on X . This first step is based on the so called *heat-flow method*. In the second step, the authors show how to find functions F_j and G_j with the desired properties by employing a so-called *Bellman function*. Their third step consists in establishing the inequality (1.1) by reducing the problem to the case that $A = I - E_\lambda$ on $\mathbb{C}^2$, where

$$E_\lambda = \begin{pmatrix} 0 & \overline{\lambda} \\ \lambda & 0 \end{pmatrix}, \quad (\lambda \in \mathbb{T}).$$

The underlying reduction procedure is actually well-known in the literature, but has been used mainly for symmetric *sub-Markovian* semigroups, i.e., under the additional assumption that all $S_t \geq 0$. Here, the last step becomes considerably simpler, since then one need only consider the cases $A = I - E_1$ and $A = I$.

One intention with the present paper is to look more carefully at the employed reduction techniques (Section 3) and prove a general theorem (Theorem 2.2) that puts the abovementioned “third step” on a formal basis. Where the authors of [2] confine their arguments to their specific case of Bellman functions, here we treat general functions F_j and G_j and hence pave the way for further applications.

It turns out that the heart of the matter are results about representing bilinear forms $(f, g) \mapsto \int_L Tf \cdot g \, d\mu$ as integrals over product spaces like

$$\int_L Tf \cdot g \, d\mu = \int_{K \times L} f(x)g(y) \, d\mu_T(x, y).$$

(Here, K and L are compact spaces, μ is a positive regular Borel measure on L and $T : C(K) \rightarrow L^1(L, \mu)$ is a linear operator.) These results go back to Grothendieck’s work on tensor products and “integral” bilinear forms [9]. They are “well-known” in the sense that they could — on a careful reading — be obtained from standard texts on tensor products and Banach lattices, such as [21, Chap. IV]. However, it seems that the communities of those people who are familiar with these facts in their abstract form and those who would like to apply them to more concrete situations are almost disjoint. Our exposition, forming the contents of Section 4, can thus be viewed as an attempt to increase the intersection of these two communities.

After this excursion into abstract operator theory, in Section 5 we turn back to the proof of Theorem 2.2. Then, as an application, we consider the

question about the optimal *angle of analyticity* on L^p of a symmetric contraction semigroup $(S_t)_{t \geq 0}$. For the sub-Markovian case this question has been answered long ago, in fact, by the very methods which we just mentioned and which form the core content of this paper. The general symmetric case has only recently been settled by Kriegler in [16]. Kriegler's proof rests on arguments from non-commutative operator theory, but Carbonaro and Dragičević show in [2] that the result can also be derived as a corollary from their results involving Bellman-functions. We shall point out in Section 6 below that the Bellman function of Carbonaro and Dragičević is not really needed here, and that one can prove the general case by essentially the same arguments as used in the sub-Markovian case.

Terminology and Notation. In this paper, $X := (X, \Sigma, \mu)$ denotes a general measure space. (Sometimes we shall suppose in addition that μ is a finite measure, but we shall always make this explicit.) Integration with respect to μ is abbreviated by

$$\int_X f := \int_X f \, d\mu$$

whenever it is convenient. The corresponding L^p -space for $0 < p \leq \infty$ is denoted by $L^p(X)$, but if the underlying measure space is understood, we shall simply write L^p . Whenever $1 \leq p \leq \infty$ is fixed we denote by q the *dual exponent*, i.e., the unique number $q \in [1, \infty]$ such that $\frac{1}{p} + \frac{1}{q} = 1$.

With the symbol $\mathcal{M}(X; \mathbb{C}^d)$ ($\mathcal{M}(X)$ in the case $d=1$) we denote the space of $\mathbb{C}^d$ -valued measurable functions on X , modulo equality almost everywhere. We shall tacitly identify $\mathcal{M}(X; \mathbb{C}^d)$ with $\mathcal{M}(X)^d$ and use the notation

$$\mathbf{f} = (f_1, \dots, f_d)$$

to denote functions into $\mathbb{C}^d$. For a set $M \subseteq \mathbb{C}^d$ and $\mathbf{f} = (f_1, \dots, f_d) \in \mathcal{M}(X; \mathbb{C}^d)$ as above, we write “ $(f_1, \dots, f_d) \in M$ almost everywhere” shorthand for: “ $(f_1(x), \dots, f_d(x)) \in M$ for μ -almost all $x \in X$.” By abuse of notation, if $F : \mathbb{C}^d \rightarrow \mathbb{C}$ is measurable and $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ we write $F(\mathbf{f})$ to denote the function $F \circ \mathbf{f}$, i.e., $F(\mathbf{f})(x) = F(f_1(x), \dots, f_d(x))$.

The letters $K, L, \dots$ usually denote compact and sometimes locally compact Hausdorff spaces. We abbreviate this by simply saying that $K, L, \dots$ are (locally) compact. If K is locally compact, then $C_c(K)$ denote the space of continuous functions on K with compact support, and $C_0(K)$ is the sup-norm closure of $C_c(K)$ within the Banach space of all bounded continuous functions. If K is compact, then of course $C_c(K) = C_0(K) = C(K)$.

If K is (locally) compact then, by the Riesz representation theorem, the dual space of $C(K)$ ($C_0(K)$) is isometrically and lattice isomorphic to $M(K)$, the space of complex regular Borel measures on K , with the total variation (norm) as absolute value (norm). A (locally) compact measure space is a pair (K, ν) where K is (locally) compact and ν is a positive regular Borel measure on K . (If K is locally compact, the measure ν need not be finite.)

We work with complex Banach spaces by default. In particular, L^p -spaces have to be understood as consisting of complex-valued functions. For

an operator T with domain and range being spaces of complex-valued functions, the *conjugate operator* is defined by $\overline{T}f := \overline{Tf}$, and the *real part* and *imaginary part* are defined by

$$\operatorname{Re} T := \frac{1}{2}(T + \overline{T}) \quad \text{and} \quad \operatorname{Im} T := \frac{1}{2i}(T - \overline{T}),$$

respectively. For Banach spaces E and F we use the symbol $\mathcal{L}(E; F)$ to denote the space of bounded linear operators from E to F and $E' = \mathcal{L}(E; \mathbb{C})$ for the dual space. The dual of an operator $T \in \mathcal{L}(E; F)$ is denoted by $T' \in \mathcal{L}(F', E')$.

If K is locally compact, $X = (X, \Sigma, \mu)$ is a measure space and $T : C_c(K) \rightarrow L^1(X)$ is a linear operator, then $T'\mu$ denotes the linear functional on $C_c(K)$ defined by

$$\langle f, T'\mu \rangle := \int_X Tf \, d\mu \quad (f \in C_c(K)).$$

If T is bounded for the uniform norm on $C_c(K)$ then $T'\mu$ is bounded too, and we identify it with the complex regular Borel measure in $M(K)$. If T is not bounded but positive, then, again by the Riesz representation theorem, $T'\mu$ can be identified with a positive (but infinite) regular Borel measure on K .

At some places we use some basic notions of Banach lattice theory (e.g., lattice homomorphism, ideal, order completeness). The reader unfamiliar with this terminology can consult [5, Chap. 7] for a brief account. However, the only Banach lattice that appears here and is not a function space will be $M(K)$, where K is locally compact.

2. Main Results

An **absolute contraction**, or a **Dunford–Schwartz operator**, over a measure space X is an operator $T : L^1 \cap L^\infty \rightarrow L^1 + L^\infty$ satisfying $\|Tf\|_p \leq \|f\|_p$ for $p = 1$ and $p = \infty$. It is then well-known that T extends uniquely and consistently to linear contraction operators $T_p : L^p \rightarrow L^p$ for $1 \leq p < \infty$, and $T_\infty : L^{(\infty)} \rightarrow L^{(\infty)}$, where $L^{(\infty)}$ is the closed linear hull of $L^1 \cap L^\infty$ within L^∞ . It is common to use the single symbol T for each of the operators T_p .

An absolute contraction T is **sub-Markovian** if it is **positive**, i.e., if $Tf \geq 0$ whenever $f \geq 0$, $f \in L^1 \cap L^\infty$. (Then also its canonical extension T_p to L^p , $1 \leq p < \infty$ and $L^{(\infty)}$, $p = \infty$, is positive.) This terminology is coherent with [20, Def. 2.12].

An absolute contraction T is called **Markovian**, if it satisfies

$$f \leq b\mathbf{1} \quad \implies \quad Tf \leq b\mathbf{1}$$

for every $b \in \mathbb{R}$ and $f \in L^1 \cap L^\infty$. (Here, $\mathbf{1}$ is the constant function with value equal to 1.) In particular, T is positive, i.e., sub-Markovian. If the measure space X is finite, an absolute contraction is Markovian if and only if T is positive and $T\mathbf{1} = \mathbf{1}$. This is easy to see, cf. [10, Lemma 3.2].

An operator $T : L^1 \cap L^\infty \rightarrow L^1 + L^\infty$ is **symmetric** if

$$\int_X Tf \cdot \bar{g} = \int_X f \cdot \overline{Tg}$$

for all $f, g \in L^1 \cap L^\infty$. A symmetric operator is an absolute contraction if and only if it is L^∞ -contractive if and only if it is L^1 -contractive; and in this case the canonical extension to L^2 is a bounded self-adjoint operator.

A (strongly continuous) **absolute contraction semigroup** over X is a family $(S_t)_{t \geq 0}$ of absolute contractions on X such that $S_0 = I$, $S_{t+s} = S_t S_s$ for all $t, s \geq 0$ and

$$\|f - S_t f\|_p \rightarrow 0 \quad \text{as } t \searrow 0 \quad (2.1)$$

for all $f \in L^1 \cap L^\infty$ and all $1 \leq p < \infty$. It follows that the operator family $(S_t)_{t \geq 0}$ can be considered a strongly continuous semigroup on each space L^p , $1 \leq p < \infty$. *We shall always assume this continuity property even when it is not explicitly mentioned.* An absolute contraction semigroup $(S_t)_{t \geq 0}$ is called a **symmetric contraction semigroup** (**symmetric (sub-)Markovian semigroup**) if each operator S_t , $t \geq 0$, is symmetric (symmetric and (sub-)Markovian).

Remarks 2.1. 1) A symmetric sub-Markovian semigroup is called a “symmetric diffusion semigroup” in the classical text [23]. It appears that the “diffusion semigroups” of operator space theory [16, Def. 2] lack the property of positivity, and hence do not specialize to Stein’s concept in the commutative case, but rather to what we call “symmetric contraction semigroups” here.

2) As Voigt [25] has shown, the strong continuity assumption (2.1) for $p \neq 2$ is a consequence of the case $p = 2$ together with the requirement that all operators S_t are L^p -contractions.

Given an absolute contraction semigroup $(S_t)_{t \geq 0}$ one can consider, for $1 \leq p < \infty$, the negative generator $-A_p$ of the strongly continuous semigroup $(S_t)_{t \geq 0}$ on L^p , defined by

$$\begin{aligned} \text{dom}(A_p) &= \{f \in L^p : \lim_{t \searrow 0} \frac{1}{t}(f - S_t f) \text{ exists in } L^p\}, \\ A_p f &= \lim_{t \searrow 0} \frac{1}{t}(f - S_t f). \end{aligned}$$

The operators A_p are compatible for different indices p , a fact which is easily seen by looking at the resolvent of A_p

$$(I + A_p)^{-1} f = \int_0^\infty e^{-t} S_t f \, dt \quad (f \in L^p, 1 \leq p < \infty).$$

Hence, it is reasonable to drop the index p and simply write A instead of A_p .

In order to formulate the main result, we first look at the very special case that the underlying measure space consists of two atoms with equal mass. Let this (probability) space be denoted by Z_2 , i.e.,

$$Z_2 := (\{0, 1\}, 2^{\{0, 1\}}, \zeta_2).$$

Then, for $1 \leq p < \infty$, $L^p(Z_2) = \mathbb{C}^2$ with norm

$$\left\| \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \right\|_p^p = \frac{1}{2}(|z_1|^p + |z_2|^p).$$

The scalar product on the Hilbert space $H = L^2(Z_2)$ is

$$\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \cdot_{Z_2} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \frac{1}{2}(z_1 \overline{w_1} + z_2 \overline{w_2}).$$

Symmetric operators on $L^2(Z_2)$ are represented by matrices

$$T = \begin{pmatrix} a & \overline{w} \\ w & b \end{pmatrix}$$

with $a, b \in \mathbb{R}$. The property that T is an absolute contraction is equivalent with the conditions $|a| + |w| \leq 1$ and $|b| + |w| \leq 1$. Thus, the absolute contractions on Z_2 form a closed convex set

$$C_2 := \left\{ \begin{pmatrix} a & \overline{w} \\ w & b \end{pmatrix} \mid a, b \in \mathbb{R}, w \in \mathbb{C}, \max\{|a|, |b|\} \leq 1 - |w| \right\},$$

and it is easy to see that each matrix

$$E_\lambda := \begin{pmatrix} 0 & \overline{\lambda} \\ \lambda & 0 \end{pmatrix}, \quad \lambda \in \mathbb{T},$$

is an extreme point of C_2 . We can now formulate the desired (meta-)theorem.

Theorem 2.2 (Symmetric Contraction Semigroups). *Let $m, d \in \mathbb{N}$, $1 \leq p < \infty$ and let, for each $1 \leq j \leq m$, $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ be measurable functions. For any generator $-A$ of a symmetric contraction semigroup over a measure space X consider the following statement:*

“For all measurable functions $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in \text{dom}(A_p)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \text{Re} \int_X A F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \geq 0.”$$

Then this statement holds true provided it holds true whenever X is replaced by Z_2 and A is replaced by $I - E_\lambda$, $\lambda \in \mathbb{T}$.

If, in addition, the semigroup is sub-Markovian, we have an even better result. In slightly different form (but with with more or less the same method), this result has been obtained by Huang in [12, Theorem 2.2].

Theorem 2.3 (Sub-Markovian Semigroups). *Let $m, d \in \mathbb{N}$, $1 \leq p < \infty$ and let, for each $1 \leq j \leq m$, $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ be measurable functions. For any generator $-A$ of a symmetric sub-Markovian semigroup over a measure space X consider the following statement:*

“For all measurable functions $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in \text{dom}(A_p)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \text{Re} \int_X A F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \geq 0.”$$

Then this statement holds true provided it holds true whenever X is replaced by Z_2 and A is replaced by $I - E_1$ and by I .

The second condition here (that the statement holds for Z_2 and $A = I$) just means that the scalar inequality

$$\sum_{j=1}^m \operatorname{Re} F_j(x) G_j(x) \geq 0$$

holds for all $x \in \mathbb{C}^d$, cf. Lemma 5.1 below.

Finally, we suppose that the measure space X is finite and the semigroup is *Markovian*, i.e., $S_t \geq 0$ and $S_t \mathbf{1} = \mathbf{1}$ for each $t \geq 0$. Then we have an even simpler criterion.

Theorem 2.4 (Markovian Semigroups). *Let $m, d \in \mathbb{N}$, $1 \leq p < \infty$ and let, for each $1 \leq j \leq m$, $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ be measurable functions. For any generator $-A$ of a symmetric Markovian semigroup over a measure space X consider the following statement:*

“For all measurable functions $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in \operatorname{dom}(A_p)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \operatorname{Re} \int_X A F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \geq 0.”$$

Then this statement holds true provided it holds true whenever X is replaced by Z_2 and A is replaced by $I - E_1$.

The proofs of Theorems 2.2–2.4 are completed in Section 5 below after we have performed some preparatory reductions (Section 3) and provided some results from abstract operator theory (Section 4).

3. Reduction Steps

In this section we shall formulate and prove three results that, when combined, reduce the proof of Theorem 2.2 to the case when $X = (K, \mu)$ is a compact measure space, μ has full support, $L^\infty(X) = C(K)$, and $A = I - T$, where T is a single symmetric absolute contraction on X . These steps are, of course, well-known, but for the convenience of the reader we discuss them in some detail.

3.1. Reduction to Bounded Operators

Suppose that $(S_t)_{t \geq 0}$ is an absolute contraction semigroup on X with generator $-A$. Then each operator $-(I - S_\varepsilon)$ is itself the (bounded) generator of a (uniformly continuous) absolute contraction semigroup $(e^{-t(I - S_\varepsilon)})_{t \geq 0}$ on X . By definition of A ,

$$\frac{1}{\varepsilon} (I - S_\varepsilon) g \rightarrow A g \quad \text{as } \varepsilon \searrow 0$$

in L^p for $g \in \operatorname{dom}(A_p)$. We thus have the following first reduction result.

Proposition 3.1. *Let $m, d \in \mathbb{N}$, $1 \leq p < \infty$ and let, for each $1 \leq j \leq m$, $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ be measurable functions. For any generator $-A$ of an absolute contraction semigroup $(S_t)_{t \geq 0}$ over a measure space X consider the following statement:*

“For all measurable functions $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in \text{dom}(A_p)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \text{Re} \int_X A F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \geq 0.”$$

Then this statement holds true provided it holds true whenever A is replaced by $I - S_\varepsilon$, $\varepsilon > 0$.

Note that in the case $A = I - T$, the condition $F_j(\mathbf{f}) \in \text{dom}(A_p)$ just asserts that $F_j(\mathbf{f}) \in L^p$.

3.2. Reduction to a Finite Measure Space

Now it is shown that one may confine to finite measure spaces. For a given measure space $X = (X, \Sigma, \mu)$, the set

$$\Sigma_{\text{fin}} := \{B \in \Sigma : \mu(B) < \infty\}$$

is directed with respect to set inclusion. For asymptotic statements with respect to this directed set we use the abbreviation “ $B \rightarrow X$ ”. The multiplication operators

$$M_B : \mathcal{M}(X; \mathbb{C}^d) \rightarrow \mathcal{M}(X; \mathbb{C}^d), \quad M_B \mathbf{f} := \mathbf{1}_B \cdot \mathbf{f}$$

form a net, with $M_B \rightarrow I$ strongly on L^p as $B \rightarrow X$ and $1 \leq p < \infty$. It follows that for a given absolute contraction T on X and functions $f \in L^p(X)$ and $g \in L^q(X)$

$$\int_X (I - T) M_B f \cdot (M_B g) \rightarrow \int_X (I - T) f \cdot g \quad \text{as } B \rightarrow X.$$

For given $B \in \Sigma_{\text{fin}}$ we form the finite measure space (B, Σ_B, μ_B) , where $\Sigma_B := \{C \in \Sigma : C \subseteq B\}$ and $\mu_B := \mu|_{\Sigma_B}$. Then we have the extension operator

$$\text{Ext}_B : \mathcal{M}(B; \mathbb{C}^d) \rightarrow \mathcal{M}(X; \mathbb{C}^d), \quad \text{Ext}_B \mathbf{f} = \begin{cases} \mathbf{f} & \text{on } B \\ 0 & \text{on } X \setminus B, \end{cases}$$

and the restriction operator

$$\text{Res}_B : \mathcal{M}(X; \mathbb{C}^d) \rightarrow \mathcal{M}(B; \mathbb{C}^d), \quad \text{Res}_B f := f|_B.$$

Note that $\text{Ext}_B \text{Res}_B = M_B$ and $\text{Res}_B \text{Ext}_B = I$ and

$$\int_B \text{Res}_B f \, d\mu_B = \int_X M_B f \, d\mu \quad (f \in L^1(X)).$$

A short computation yields that $\text{Res}_B^* = \text{Ext}_B$ between the respective L^2 -spaces. Hence, if T is a (symmetric) absolute contraction on $X = (X, \Sigma, \mu)$, then the operator

$$T_B := \text{Res}_B T \text{Ext}_B$$

is a (symmetric) absolute contraction on (B, Σ_B, μ_B) . Another short computation reveals that

$$\int_X (I - T)M_B f \cdot (M_B g) d\mu = \int_B (I_{L^p(B)} - T_B)(\text{Res}_B f) \cdot (\text{Res}_B g) d\mu_B$$

whenever $f \in L^p(X)$ and $g \in L^q(X)$. Finally, suppose that $F : \mathbb{C}^d \rightarrow \mathbb{C}$ is measurable and suppose that $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ is such that $F(\mathbf{f}) \in L^p(X)$. Then

$$\text{Res}_B[F(\mathbf{f})] = F(\text{Res}_B \mathbf{f}) \in L^p(B).$$

Combining all these facts yields our second reduction result.

Proposition 3.2. *Let $m, d \in \mathbb{N}$, $1 \leq p < \infty$ and let, for each $1 \leq j \leq m$, $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ be measurable functions. For any absolute contraction T over a measure space $X = (X, \Sigma, \mu)$ consider the following statement:*
“For all measurable functions $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in L^p(X)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \text{Re} \int_X (I - T)F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \geq 0.”$$

Then this statement holds true provided it holds true whenever $X = (X, \Sigma, \mu)$ is replaced by (B, Σ_B, μ_B) and T is replaced by T_B , where $B \in \Sigma_{\text{fin}}$.

Finally, we observe that if T is sub-Markovian (=positive) or Markovian, then so is each of the operators $T_B = \text{Res}_B T \text{Ext}_B$, $B \in \Sigma_{\text{fin}}$.

3.3. Reduction to a Compact Measure Space

In the next step we pass from general finite measure spaces to *compact* spaces with a finite positive Borel measure on it.

Let $X = (X, \Sigma, \mu)$ be a *finite* measure space. The space $L^\infty(X)$ is a commutative, unital C^* -algebra, whence by the Gelfand–Naimark theorem there is a compact space K , the Gelfand space, and an isomorphism of unital C^* -algebras

$$\Phi : L^\infty(X) \rightarrow C(K).$$

In particular, Φ is an isometry. Since the order structure is determined by the C^* -algebra structure (an element f is ≥ 0 if and only if there is g such that $f = g\bar{g}$), Φ is also an isomorphism of complex Banach lattices. The following auxiliary result is, essentially, a consequence of the Stone–Weierstrass theorem.

Lemma 3.3. *In the situation from above, let $M \subseteq \mathbb{C}^d$ be compact and let $f_1, \dots, f_d \in L^\infty(X)$ be such that $(f_1, \dots, f_d) \in M$ μ -almost everywhere. Then $(\Phi f_1, \dots, \Phi f_d) \in M$ everywhere on K and*

$$\Phi(F(f_1, \dots, f_d)) = F(\Phi f_1, \dots, \Phi f_d) \tag{3.1}$$

for all continuous functions $F \in C(M)$.

Proof. Suppose first that $M = B[0, r] := \{x \in \mathbb{C}^d : \|x\|_\infty \leq r\}$ for some $r > 0$. Then the condition “ $(f_1, \dots, f_d) \in M$ almost everywhere” translates into the inequalities $|f_j| \leq r\mathbf{1}$ (almost everywhere) for all $j = 1, \dots, d$, and hence one has also $|\Phi f_j| \leq r\Phi\mathbf{1} = r\mathbf{1}$ (pointwise everywhere) for all $j = 1, \dots, d$. It follows that $F(\Phi f_1, \dots, \Phi f_d)$ is well-defined.

Now, the set of functions $F \in C(M)$ such that (3.1) holds is a closed conjugation-invariant subalgebra of $C(M)$ that separates the points and contains the constants. Hence, by the Stone-Weierstrass theorem, it is all of $C(M)$.

For general M one can proceed in the same way provided one can assure that $(\Phi f_1, \dots, \Phi f_d) \in M$ everywhere on K . Let $y \in \mathbb{C}^d \setminus M$ and let F be any continuous function with compact support on $\mathbb{C}^d$ such that $F = 0$ on M and $F(y) = 1$. Let $r > 0$ be so large that $M \subseteq B[0, r]$ and consider F as a function on $B[0, r]$. Then $0 = \Phi(0) = \Phi(F(f_1, \dots, f_d)) = F(\Phi f_1, \dots, \Phi f_d)$, whence y cannot be in the image of $(\Phi f_1, \dots, \Phi f_d)$. $\square$

By the Riesz-Markov representation theorem, there is a unique regular Borel measure ν on K such that

$$\int_X f = \int_K \Phi f \, d\nu$$

for all $f \in L^\infty(X)$. It follows from Lemma 3.3 that $|\Phi f|^p = \Phi(|f|^p)$ for every $1 \leq p < \infty$ and every $f \in L^\infty(X)$. Therefore, Φ is an isometry with respect to each p -norm. It follows that Φ extends to an isometric (lattice) isomorphism

$$\Phi : L^1(X) \rightarrow L^1(K, \nu).$$

It is shown in Appendix A that Φ , furthermore, extends canonically (and uniquely) to a unital $*$ -algebra and lattice isomorphism

$$\Phi : \mathcal{M}(X) \rightarrow \mathcal{M}(K, \nu).$$

The compact measure space (K, ν) (together with the mapping Φ) is called the **Stone model** of the probability space X . Note that under the lattice isomorphism Φ the respective L^∞ -spaces must correspond to each other, whence it follows that $L^\infty(K, \mu) = C(K)$ in the obvious sense.

We use the canonical extension to vector-valued functions $\Phi : \mathcal{M}(X; \mathbb{C}^d) \rightarrow \mathcal{M}(K, \nu; \mathbb{C}^d)$ of the Stone model. By Theorem A.3,

$$\Phi(F(\mathbf{f})) = F(\Phi \mathbf{f}) \quad \nu\text{-almost everywhere}$$

for all measurable functions $\mathbf{f} = (f_1, \dots, f_d) \in \mathcal{M}(X; \mathbb{C}^d)$ and all measurable functions $F : \mathbb{C}^d \rightarrow \mathbb{C}$. Hence, we arrive at the next reduction result.

Proposition 3.4. *Let $m, d \in \mathbb{N}$, $1 \leq p < \infty$ and let, for each $1 \leq j \leq m$, $F_j, G_j : \mathbb{C}^d \rightarrow \mathbb{C}$ be measurable functions. For any absolute contraction T over a probability space X consider the following statement:*

“For all measurable functions $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in L^p(X)$ and $G_j(\mathbf{f}) \in L^q(X)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \operatorname{Re} \int_X [(I - T)F_j(\mathbf{f})] \cdot G_j(\mathbf{f}) \geq 0.”$$

Then this statement holds true provided it holds true if X is replaced by (K, ν) and T is replaced by $\Phi T \Phi^{-1}$, where (K, ν) and

$$\Phi : \mathcal{M}(X) \rightarrow \mathcal{M}(K, \nu)$$

is the Stone model of X .

As in the reduction step before, we observe that the properties of being symmetric, sub-Markovian or Markovian are preserved during the reduction process, i.e., in passing from T to $\Phi^{-1}T\Phi$.

Remark 3.5. In the late 1930’s and beginning 1940’s, several representation results for abstract structures were developed first by Stone [24] (for Boolean algebras), then by Gelfand [7, 8] (for normed algebras) and Kakutani [13, 14] (for AM - and AL -spaces). However, it is hard to determine when for the first time there was made effective use of these results in a context similar to ours. Halmos in his paper [11] on a theorem of Dieudonné on measure disintegration employs the idea but uses Stone’s original theorem. A couple of years later, Segal [22, Thm. 5.4] revisits Dieudonné’s theorem and gives a proof based on algebra representations. (He does not mention Gelfand–Naimark, but only says “by well-known results”.)

In our context, the idea — now through the Gelfand–Naimark theorem — was employed by Nagel and Voigt [19] in order to simplify arguments in the proof of Liskevich and Perelmutter [17] on the optimal angle of analyticity in the sub-Markovian case, see Section 6 below. Through Ouhabaz’ book [20] it has become widely known in the field, and also Carbonaro and Dragičević [2, p.19] use this idea.

4. Operator Theory

In order to proceed with the proof of the main theorem (Theorem 2.2) we need to provide some results from the theory of operators of the form $T : C(K) \rightarrow L^1(L, \mu)$, where K and L are compact.¹ For the application to symmetric contraction semigroups as considered in the previous sections, we only need the case that $C(L) = L^\infty(L, \mu)$, and this indeed would render simpler some of the proofs below. However, a restriction to this case is artificial, and we develop the operator theory in reasonable generality.

¹The case that K and L are locally compact is touched upon in some additional remarks.

4.1. The Linear Modulus

In this section we introduce the linear modulus of an order-bounded operator $T : C(K) \rightarrow L^1(X)$. This can be treated in the framework of general Banach lattices, see [21, Chapter IV, §1], but due to our concrete situation, things are a little easier than in an abstract setting.

Let $X = (X, \Sigma, \mu)$ be a measure space and let K be compact. A linear operator $T : C(K) \rightarrow L^1(X)$ is called **order-bounded** if for each $0 \leq f \in C(K)$ there is $0 \leq h \in L^1(X)$ such that

$$|Tu| \leq h \quad \text{for all } u \in C(K) \text{ with } |u| \leq f.$$

And T is called **regular** if it is a linear combination of positive operators. It is clear that each regular operator is order-bounded. The converse also holds, by the following construction.

Suppose that $T : C(K) \rightarrow L^1(X)$ is order-bounded. Then, for $0 \leq f \in C(K)$ let

$$|T|f := \sup\{|Tg| : g \in C(K), |g| \leq f\} \quad (4.1)$$

as a supremum in the lattice sense. (This supremum exists since the set on the right hand side is order bounded by hypothesis and L^1 is order complete, see [5, Cor. 7.8].)

Lemma 4.1. *Suppose that $T : C(K) \rightarrow L^1(X)$ is order-bounded. Then the mapping $|T|$ defined by (4.1) extends uniquely to a positive operator*

$$|T| : C(K) \rightarrow L^1(X).$$

Moreover, the following assertions hold:

- a) $|Tf| \leq |T||f|$ for all $f \in C(K)$.
- b) $\|T\| \leq \||T|\|$,
- c) $\overline{T}$ is order-bounded and $|\overline{T}| = |T|$.
- d) If $S : C(K) \rightarrow L^1(X)$ is order-bounded, then $S+T$ is also order-bounded, and $|S+T| \leq |S| + |T|$.

The operator $|T| : C(K) \rightarrow L^1(X)$ whose existence is asserted in the theorem is called the **linear modulus** of T .

Proof. For the first assertion, it suffices to show that $|T|$ is additive and positively homogeneous. The latter is straightforward, so consider additivity. Fix $0 \leq f, g \in C(K)$ and let $u \in C(K)$ with $|u| \leq f + g$. Define

$$u_1 = \frac{fu}{f+g}, \quad u_2 = \frac{gu}{f+g},$$

where $u_1 = u_2 = 0$ on the set $[f+g=0]$. Then $u_1, u_2 \in C(K)$, $|u_1| \leq f$, $|u_2| \leq g$ and $u_1 + u_2 = u$. Hence

$$|Tu| \leq |Tu_1| + |Tu_2| \leq |T|f + |T|g$$

and taking the supremum with respect to u we obtain $|T|(f+g) \leq |T|f + |T|g$. Conversely, let $u, v \in C(K)$ with $|u| \leq f$ and $|v| \leq g$. Then, for any $\alpha \in \mathbb{C}^2$ with $|\alpha_1| + |\alpha_2| \leq 1$ we have $|\alpha_1 u + \alpha_2 v| \leq f + g$, and hence

$$|Tu| + |Tv| = \sup_{\alpha} |\alpha_1 Tu + \alpha_2 Tv| = \sup_{\alpha} |T(\alpha_1 u + \alpha_2 v)| \leq |T|(f + g).$$

Taking suprema with respect to u and v we arrive at $|T|f + |T|g \leq |T|(f + g)$. The remaining statements are now easy to establish. $\square$

Suppose that $T : C(K) \rightarrow L^1(X)$ is order-bounded, so that $|T|$ exists. Then, by Lemma 4.1, also $\operatorname{Re} T$ and $\operatorname{Im} T$ are order-bounded. If T is real, i.e., if $T = \overline{T}$, then clearly $T \leq |T|$, and hence $T = |T| - (|T| - T)$ is regular. It follows that every order-bounded operator is regular. (See also [21, IV.1, Prop.s 1.2. and 1.6].)

Let us turn to another characterization of order-boundedness. If $T : C(K) \rightarrow L^1(X)$ is order-bounded and $|T|$ is its linear modulus, we denote by $|T|'\mu$ the unique element $\nu \in M(K)$ such that

$$\int_K f d\nu = \int_X |T|f \quad \text{for all } f \in C(K).$$

It is then easy to see that T extends to a contraction $T : L^1(K, \nu) \rightarrow L^1(X)$. We shall see that the existence of a positive regular Borel measure ν on K with this property characterizes the order-boundedness. The key is the following general result, which has (probably) been established first by Grothendieck [9, p.67, Corollaire].

Lemma 4.2. *Let X, Y be measure spaces and let $T : L^1(Y) \rightarrow L^1(X)$ be a bounded operator. Then for any finite sequence $f_1, \dots, f_n \in L^1(Y)$*

$$\int_X \sup_{1 \leq j \leq n} |Tf_j| \leq \|T\| \int_Y \sup_{1 \leq j \leq n} |f_j|.$$

Proof. By approximation, we may suppose that all the functions f_j are integrable step functions with respect to one finite partition $(A_k)_k$. We use the variational form

$$\sup_{1 \leq j \leq n} |z_j| = \sup \left\{ \left| \sum_j^n \alpha_j z_j \right| : \alpha \in \ell_n^1, \|\alpha\|_1 \leq 1 \right\}$$

for complex numbers $z_1, \dots, z_n$. Then, with $f_j = \sum_k c_{jk} \mathbf{1}_{A_k}$,

$$\begin{aligned} \sup_{1 \leq j \leq n} |Tf_j| &= \sup_{\alpha} \left| \sum_j^n \sum_k \alpha_j c_{jk} T\mathbf{1}_{A_k} \right| \\ &\leq \sup_{\alpha} \sum_k \|\alpha\|_1 \left(\sup_{1 \leq j \leq n} |c_{jk}| \right) |T\mathbf{1}_{A_k}| \\ &= \sum_k \left(\sup_{1 \leq j \leq n} |c_{jk}| \right) |T\mathbf{1}_{A_k}|. \end{aligned}$$

Integrating yields

$$\int_X \sup_{1 \leq j \leq n} |Tf_j| \leq \sum_k \left(\sup_{1 \leq j \leq n} |c_{jk}| \right) \|T\mathbf{1}_{A_k}\|_1$$

$$\begin{aligned}
&\leq \|T\| \sum_k \left(\sup_{1 \leq j \leq n} |c_{jk}| \right) \|\mathbf{1}_{A_k}\|_1 \\
&= \|T\| \int_Y \sum_k \left(\sup_{1 \leq j \leq n} |c_{jk}| \right) \mathbf{1}_{A_k} = \|T\| \int_Y \sup_{1 \leq j \leq n} |f_j|. \quad \square
\end{aligned}$$

We can now formulate the main result of this section.

Theorem 4.3. *Let $X = (X, \Sigma, \mu)$ be any measure space and $T : C(K) \rightarrow L^1(X)$ a linear operator. Then the following assertions are equivalent:*

- (i) *T is order-bounded.*
- (ii) *T is regular.*
- (iii) *There is a positive regular Borel measure $\nu \in M(K)$ such that T extends to a contraction $L^1(K, \nu) \rightarrow L^1(X)$.*

If (i)–(iii) hold, then

$$|T|' \mu = \min \{ \nu \in M_+(K) : \|Tf\|_{L^1(X)} \leq \|f\|_{L^1(K, \nu)} \text{ for all } f \in C(K) \}.$$

In particular, if $0 \leq \nu \in M(K)$ is such that T extends to a contraction $L^1(K, \nu) \rightarrow L^1(X)$, then so does $|T|$.

Proof. The implications (i) $\Leftrightarrow$ (ii) $\Rightarrow$ (iii) have already been established. Moreover, if (i) holds then it follows from the inequality $|Tf| \leq |T||f|$ that $\|Tf\|_1 \leq \|f\|_{L^1(K, \nu)}$ with $\nu = |T|' \mu$.

On the other hand, suppose (iii) holds and that $0 \leq \nu \in M(K)$ is such that $\int_X |Tf| \leq \int_K |f| d\nu$ for all $f \in C(K)$. Let $0 \leq f \in C(K)$, $n \in \mathbb{N}$ and $u_j \in C(K)$ with $|u_j| \leq f$ ($1 \leq j \leq n$). Then, by Lemma 4.2,

$$\int_X \sup_{1 \leq j \leq n} |Tu_j| \leq \int_K \sup_{1 \leq j \leq n} |u_j| d\nu \leq \int_K f d\nu.$$

Now, any upwards directed and norm bounded net in L^1_+ is order-bounded and converges in L^1 -norm towards its supremum, see [5, Thm. 7.6]. It follows that T is order-bounded, and

$$\int_X |T|f \leq \int_K f d\nu.$$

Consequently, $|T|' \mu \leq \nu$, as claimed. $\square$

Remarks 4.4. 1) Suppose that (i)–(iii) of Theorem 4.3 hold. Then $|T' \mu| \leq |T|' \mu$, and equality holds if and only if T extends to a contraction $T : L^1(K, |T' \mu|) \rightarrow L^1(X)$.

2) The modulus mapping $T \mapsto |T|$ turns $\mathcal{L}^r(C(K), L^1(X))$, the set of regular operators, into a complex Banach lattice with the norm $\|T\|_r := \||T|\|$, see [21, Chap. IV, §1].

3) All the results of this section hold *mutatis mutandis* for linear operators $T : C_c(Y) \rightarrow L^1(X)$, where Y is a locally compact space and $C_c(Y)$ is the space of continuous functions on Y with compact support.

The modulus of a linear operator appears already in the seminal work of Kantorovich [15] on operators on linear ordered spaces. For operators on an L^1 -space the linear modulus was (re-)introduced in [3] by Chacon and Krengel who probably were not aware of Kantorovich's work. Later on, their construction was generalized to order-bounded operators between general Banach lattices by Luxemburg and Zaanen in [18] and then incorporated by Schaefer in his monograph [21].

The equivalence of order-bounded and regular operators is of course a standard lemma from Banach lattice theory. Lemma 4.2 is essentially equivalent to saying that every bounded operator between L^1 -spaces is order-bounded. This has been realized by Grothendieck in [9, p.66, Prop. 10]. (Our proof differs considerably from the original one.) The equivalence of (i)–(iii) in Theorem 4.3 can also be derived from combining Theorem IV.1.5 and Corollary 1 of Theorem II.8.9 of [21]. However, the remaining part of Theorem 4.3 might be new.

4.2. Integral Representation of Bilinear Forms

In this section we aim for yet another characterization of order-bounded operators $T : C(K) \rightarrow L^1(X)$ in the case that $X = (L, \mu)$ is a compact measure space. We shall see that an operator T is order-bounded if, and only if, there is a (necessarily unique) complex regular Borel measure μ_T on $K \times L$ such that

$$\int_{K \times L} f \otimes g \, d\mu_T = \int_L (Tf) \cdot g \, d\mu \quad \text{for all } f \in C(K) \text{ and } g \in C(L). \quad (4.2)$$

This result goes essentially back to Grothendieck's characterization of "integral" operators in [9, p.141, Thm. 11], but we give ad hoc proofs avoiding the tensor product theory. The following simple lemma is the key result here.

Lemma 4.5. *Let K, L be compact spaces. Then, for any bounded operator $T : C(K) \rightarrow C(L)$ and any $\mu \in M(L)$ there is a unique complex regular Borel measure $\mu_T \in M(K \times L)$ such that (4.2) holds. Moreover, $\mu_T \geq 0$ whenever $\mu \geq 0$ and $T \geq 0$.*

Proof. The uniqueness is clear since $C(K) \otimes C(L)$ is dense in $C(K \times L)$. For the existence, let $S : C(K \times L) \rightarrow C(L)$ be given by composition of all of the operators in the following chain:

$$C(K \times L) \cong C(L; C(K)) \xrightarrow{T^\otimes} C(L; C(L)) \cong C(L \times L) \xrightarrow{D} C(L).$$

Here, $T^\otimes$ denotes the operator $G \mapsto T \circ G$ and D denotes the "diagonal contraction", defined by $DG(x) := G(x, x)$ for $x \in L$ and $G \in C(L \times L)$. Then $\mu_T := S'\mu$ satisfies the requirements, as a short argument reveals. $\square$

Remarks 4.6. 1) The formula (4.2) stays true for all choices of $f \in C(K)$ and g a bounded measurable function on L .

- 2) Our proof of Lemma 4.5 yields a formula for the integration of a general $F \in C(K \times L)$ with respect to μ_T :

$$\int_{K \times L} F(x, y) d\mu_T(x, y) = \int_L (TF(\cdot, y))(y) d\mu(y).$$

This means: fix $y \in L$, apply T to the function $F(\cdot, y)$ and evaluate this at y ; then integrate this function in y with respect to μ .

- 3) Compare this proof of Lemma 4.5 with the one given in [20, p.90/91].
 4) Lemma 4.5 remains valid if K and L are merely locally compact, and $C(\cdot)$ is replaced by $C_0(\cdot)$ at each occurrence.

Combining Lemma 4.5 with a Stone model leads to the desired general theorem.

Theorem 4.7. *Let K be compact, (L, μ) a compact measure space, and $T : C(K) \rightarrow L^1(L, \mu)$ a linear operator. Then the following assertions are equivalent:*

- (i) T is order-bounded.
- (ii) T is regular.
- (iii) T extends to a contraction $L^1(K, \nu) \rightarrow L^1(L, \mu)$ for some $0 \leq \nu \in M(K)$.
- (iv) There is a complex regular Borel measure $\mu_T \in M(K \times L)$ such that (4.2) holds.

In this case, μ_T from (iv) is unique, and if ν is as in (iii), then $|T|'\mu \leq \nu$.

Proof. It was shown in Theorem 4.3 that (i)–(iii) are pairwise equivalent.

Denote by $\pi_K : K \times L \rightarrow K$ the canonical projection. Suppose that (iv) holds and let $\nu = (\pi_K)_* |\mu_T|$, i.e.,

$$\int_K f d\nu = \int_{K \times L} f \otimes \mathbf{1} d|\mu_T| \quad (f \in C(K)).$$

Then, for $f \in C(K)$ and $g \in C(L)$ with $|g| \leq 1$,

$$\left| \int_L Tf \cdot g d\mu \right| \leq \int_{K \times L} |f| \otimes |g| d|\mu_T| \leq \int_{K \times L} |f| \otimes \mathbf{1} d|\mu_T| = \int_K |f| d\nu.$$

This implies that T extends to a contraction $L^1(K, \nu) \rightarrow L^1(L, \mu)$, whence we have (iii).

Now suppose that (i)–(iii) hold. In order to prove (iv) define the operator $S : C(K) \rightarrow L^\infty(L, \mu)$ by

$$Sf := \begin{cases} \frac{Tf}{|T|\mathbf{1}} & \text{on } [|T|\mathbf{1} > 0], \\ 0 & \text{on } [|T|\mathbf{1} = 0]. \end{cases}$$

Let $\Phi : L^1(L, \mu) \rightarrow L^1(\Omega, \tilde{\mu})$ be the Stone model of (L, μ) (see Section 3.3 above), and let us identify $L^\infty(L, \mu)$ with $C(\Omega)$ via Φ . Then $S : C(K) \rightarrow C(\Omega)$

is a positive operator. Hence we can apply Lemma 4.5 to S and the positive measure $(|T|\mathbf{1})\tilde{\mu}$ to obtain a positive measure ρ on $K \times \Omega$ such that

$$\begin{aligned} \int_{K \times \Omega} f \otimes g \, d\rho &= \int_{\Omega} Sf \cdot g \, d(|T|\mathbf{1})\tilde{\mu} = \int_{\Omega} Sf \cdot |T|\mathbf{1} \cdot g \, d\tilde{\mu} \\ &= \int_{\Omega} Tf \cdot g \, d\tilde{\mu} = \int_L Tf \cdot g \, d\mu. \end{aligned}$$

Finally, let μ_T be the pull-back of ρ to $K \times L$ via the canonical inclusion map $C(L) \rightarrow L^\infty(L, \mu) = C(\Omega)$. $\square$

Remark 4.8. With a little more effort one can extend Theorem 4.7 to the case of *locally compact* (and not necessarily finite) measure spaces (K, ν) and (L, μ) instead of compact ones, cf. Remarks 4.4 and 4.6 above. Then the decisive implication (ii) $\Rightarrow$ (iv) is proved by passing first to open and relatively compact subsets $U \subseteq K$ and $V \subseteq L$ and considering the operator $T_{U,V} : C_0(U) \rightarrow L^1(V, \mu)$. By modifying our proof, one then obtains a measure $\mu_T^{U,V}$ on $U \times V$, and finally μ_T as an inductive limit. (Of course, one has to speak of Radon measures here.) Compare this to the ad hoc approach in [6, Lemma 1.4.1].

Theorem 4.7 can also be generalized to the case that K and L are *Polish* (but not necessarily locally compact) spaces and μ is a finite positive Borel measure on L . In this case the decisive implication (ii) $\Rightarrow$ (iv) is proved as follows: first, one chooses compact metric models (K', ν') and (L', μ') for the finite Polish measure spaces (K, ν) and (L, μ) , respectively, see [5, Sec. 12.3]; by a theorem of von Neumann [5, App. F.3], the isomorphisms between the original measure spaces and their models are induced by measurable maps $\varphi : K' \rightarrow K$ and $\psi : L' \rightarrow L$, say. Theorem 4.7 yields — for the transferred operator — a representing measure on $K' \times L'$, and this is mapped by $\varphi \times \psi$ to a representing measure on $K \times L$ for the original operator.

We now combine the integral Theorem 4.7 with the construction of the modulus. We employ the notation $\pi_L : K \times L \rightarrow L$ for the canonical projection, and identify

$$L^1(L, \mu) = \{\lambda \in M(L) : |\lambda| \ll \mu\}$$

with a closed ideal in $M(L)$ via the Radon-Nikodým theorem.

Theorem 4.9. *Suppose that K and L are compact spaces and $0 \leq \mu \in M(L)$. Then, for any order-bounded operator $T : C(K) \rightarrow L^1(L, \mu)$,*

$$|\mu_T| = \mu_{|T|}.$$

The mapping

$$\mathcal{L}^r(C(K), L^1(L, \mu)) \rightarrow M(K \times L), \quad T \mapsto \mu_T$$

is an isometric lattice homomorphism onto the closed ideal

$$\{\rho \in M(K \times L) : \pi_{L*}|\rho| \in L^1(L, \mu)\}$$

of $M(K \times L)$.

Proof. It is clear that the mapping $T \mapsto \mu_T$ is linear, injective and positive. Hence $|\mu_T| \leq \mu_{|T|}$, and therefore $\pi_{L*} |\mu_T| \leq \pi_{L*} \mu_{|T|} = (|T| \mathbf{1}) \mu \in L^1(L, \mu)$. Conversely, suppose that $\rho \in M(K \times L)$ such that $\pi_{L*} |\rho| \in L^1(L, \mu)$. For $f \in C(K)$ consider the linear mapping

$$T : C(K) \rightarrow M(L), \quad (Tf)g := \int_{K \times L} f \otimes g \, d\rho.$$

Then $|Tf| \leq \|f\|_\infty \pi_{L*} |\rho|$, whence $Tf \in L^1(L, \mu)$. Hence, by construction,

$$\int_{K \times L} f \otimes g \, d\rho = \int_L Tf \cdot g \, d\mu$$

for $f \in C(K)$ and $g \in C(L)$. By Theorem 4.7, T is regular. If ρ is positive, then T is positive, too.

The proof of the converse inequality $\mu_{|T|} \leq |\mu_T|$ would now follow immediately if we used the fact (from Remark 4.4) that the modulus map turns $\mathcal{L}^r$, the set of regular operators, into a complex vector lattice. However, we want to give a different proof here.

By a standard argument, it suffices to establish the inequality

$$\int_L |T| \mathbf{1} \, d\mu \leq \int_{K \times L} \mathbf{1} \otimes \mathbf{1} \, d|\mu_T|.$$

To this end, define the positive measure ν on K by

$$\int_K f \, d\nu := \int_{K \times L} f \otimes \mathbf{1} \, d|\mu_T| \quad (f \in C(K)).$$

Given $f \in C(K)$ there is a bounded measurable function h on L such that $|Tf| = (Tf)h$ and $|h| \leq 1$. Hence,

$$\int_L |Tf| \, d\mu = \int_L Tf \cdot h \, d\mu = \int_{K \times L} f \otimes h \, d\mu_T \leq \int_{K \times L} |f| \otimes \mathbf{1} \, d\mu_T = \int_K |f| \, d\nu.$$

This means that T extends to a contraction $L^1(K, \nu) \rightarrow L^1(L, \mu)$. By Theorem 4.3, it follows that $|T|' \mu \leq \nu$, whence in particular

$$\int_L |T| \mathbf{1} \, d\mu = \int_K \mathbf{1} \, d(|T|' \mu) \leq \int_K \mathbf{1} \, d\nu = \int_{K \times L} \mathbf{1} \otimes \mathbf{1} \, d|\mu_T|.$$

This concludes the proof. $\square$

Remark 4.10. One can avoid the use of the bounded measurable function h in the second part of the proof of Theorem 4.9 by passing to the Stone model of $L^1(L, \mu)$.

In case that T has additional properties, one can extend the defining formula for the measure μ_T to some non-continuous functions.

Theorem 4.11. *Let (K, ν) and (L, μ) be compact measure spaces, and let $T : C(K) \rightarrow L^\infty(L, \mu)$ be a bounded operator that extends to a bounded operator $L^1(K, \nu) \rightarrow L^1(L, \mu)$. Then the formula*

$$\int_L Tf \cdot g \, d\mu = \int_{K \times L} f \otimes g \, d\mu_T \tag{4.3}$$

holds for all $f \in L^p(K, \nu)$, $g \in L^q(L, \mu)$ and $1 \leq p, q \leq \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. We may suppose that $T : L^1(K, \nu) \rightarrow L^1(L, \mu)$ (and hence also $|T|$) is a contraction. In a first step, we claim that the formula 4.3 holds for all bounded Baire measurable functions f, g on K, L , respectively. Indeed, this follows from a standard argument by virtue of the dominated convergence theorem and the fact that the bounded Baire-measurable functions on a compact space form the smallest set of functions that contains the continuous ones and is closed under pointwise convergence of uniformly bounded sequences, see [5, Thm. E.1].

Replacing T by $|T|$ in 4.3 we then can estimate for bounded Baire measurable functions f and g and $1 < p < \infty$

$$\begin{aligned} \int_{K \times L} |f \otimes g| \, d\mu_{|T|} &= \int_{K \times L} (|f| \otimes \mathbf{1}) \cdot (\mathbf{1} \otimes |g|) \, d\mu_{|T|} \\ &\leq \left(\int_{K \times L} |f|^p \otimes \mathbf{1} \, d\mu_{|T|} \right)^{\frac{1}{p}} \cdot \left(\int_{K \times L} \mathbf{1} \otimes |g|^q \, d\mu_{|T|} \right)^{\frac{1}{q}} \\ &= \left(\int_L |T||f|^p \, d\mu \right)^{\frac{1}{p}} \cdot \left(\int_L (|T|\mathbf{1}) \cdot |g|^q \, d\mu \right)^{\frac{1}{q}} \\ &\leq \left(\int_K |f|^p \, d\nu \right)^{\frac{1}{p}} \cdot \left(\int_L (|T|\mathbf{1}) \cdot |g|^q \, d\mu \right)^{\frac{1}{q}} = \|f\|_{L^p(\nu)} \left\| (|T|\mathbf{1})^{\frac{1}{q}} g \right\|_{L^q(\mu)}. \end{aligned}$$

It follows that if A is a ν -null Baire set of K and B is a μ -null Baire set of L , then the sets $A \times L$ and $K \times B$ are $\mu_{|T|}$ -null Baire sets of $K \times L$. Moreover, the bilinear mapping $(f, g) \mapsto f \otimes g$ extends to a bounded bilinear mapping

$$L^p(K, \nu) \times L^q(L, \mu) \rightarrow L^1(K \times L, \mu_{|T|}).$$

By interpolation, T is L^p -bounded, and hence the bilinear mapping $(f, g) \mapsto Tf \cdot g$ is a bounded bilinear mapping $L^p(K, \nu) \times L^q(L, \mu) \rightarrow L^1(L, \mu)$. Now (4.3) holds for bounded Baire-measurable functions f and g , whence by approximation for all $f \in L^p(K, \nu)$ and $g \in L^q(L, \mu)$. (Choose sequences that approximate in norm and almost everywhere.)

Finally, consider $p = 1$ (the case $q = 1$ being similar). If $g \in L^\infty(L, \mu)$ then, by choosing a Baire-measurable representative for g such that $\|g\|_\infty = \|g\|_{L^\infty(L, \mu)}$ and using the results from above, we can estimate for each $f \in L^\infty(K, \nu)$,

$$\begin{aligned} \int_{K \times L} |f \otimes g| \, d\mu_{|T|} &= \int_{K \times L} (|f| \otimes \mathbf{1}) \cdot (\mathbf{1} \otimes |g|) \, d\mu_{|T|} \\ &\leq \int_{K \times L} |f| \otimes \mathbf{1} \cdot \|g\|_\infty \, d\mu_{|T|} = \| |T| |f| \|_{L^1(L, \nu)} \|g\|_\infty \\ &\leq \|f\|_{L^1(K, \nu)} \|g\|_{L^\infty(L, \mu)}. \end{aligned}$$

The assertion then follows by approximation (almost everywhere and in norm) as before. $\square$

Remark 4.12. If an operator $T : C(K) \rightarrow L^1(L, \mu)$ factors through $L^\infty(L, \mu)$, it is of course order-bounded, and hence its modulus exists. If, in addition,

it factors even through $C(K)$, then the existence of μ_T follows from Lemma 4.5 directly and one does not have to pass through the Stone model. If (L, μ) is already its own Stone model (as is the case in the proof of Theorem 2.2 after the reduction step in Section 3.3) then also $|T|$ factors through $C(L)$, and hence Lemma 4.5 is completely sufficient to construct the measures μ_T and $\mu_{|T|}$.

Using modern tensor product terminology, we have

$$C(K \times L) = C(K) \otimes_{\varepsilon} C(L) \subseteq C(K) \otimes_{\varepsilon} L^{\infty}(L, \mu) = C(K) \otimes_{\varepsilon} L^1(L, \mu)'.$$

This implies (via the Stone model of (L, μ)) that an operator $T : C(K) \rightarrow L^1(L, \mu)$ is “integral” (in the sense of Grothendieck) if and only if there is $\mu_T \in M(K \times L)$ such that (4.2) holds. Hence, the decisive equivalence of (ii) and (iv) in Theorem 4.9 is essentially [9, p.141, Thm. 11]. Schaefer incorporates these results in his systematic study of operators between Banach lattices, see [21, IV, Theorem 5.6]. However, the property $|\mu_T| = \mu_{|T|}$, essential for our application below, does not appear there. It has been stated and proved explicitly in [2, Lemma 30], but our proof is different.

4.3. The Disintegration Theorem

In this section we develop further the results of the previous section. The endpoint will be a “disintegration” theorem for operators of the form $I - T$, where T is a symmetric absolute contraction over a compact measure space.

We start with some auxiliary results.

Proposition 4.13. *Let (K, ν) and (L, μ) be compact measure spaces and let $T : C(K) \rightarrow L^1(L, \mu)$ and $S : C(L) \rightarrow L^1(K, \nu)$ be linear operators such that*

$$\int_L T f \cdot g \, d\mu = \int_K f \cdot S g \, d\nu \quad (f \in C(K), g \in C(L)). \quad (4.4)$$

If one of the operators T and S is order-bounded, then so is the other and (4.4) holds with T and S replaced by $|T|$ and $|S|$, respectively. Moreover, $\mu_T = r_* \nu_S$, where $r : L \times K \rightarrow K \times L$ is the swapping map defined by $r(x, y) = (y, x)$.

Proof. Suppose that S is order-bounded. Then, for $f \in C(K)$ and $g \in C(L)$ with $|g| \leq 1$,

$$\left| \int_L T f \cdot g \, d\mu \right| \leq \int_K |f| \cdot |S g| \, d\nu \leq \int_K |f| (|S| \mathbf{1}) \, d\nu.$$

Hence, T extends to a contraction $L^1(K, (|S| \mathbf{1}) \nu) \rightarrow L^1(L, \mu)$, whence, by Theorem 4.3, is order-bounded and $|T|' \mu \leq (|S| \mathbf{1}) \nu$. (Recall that the unit ball of $C(L)$ is L^1 -dense in the unit ball of $L^{\infty}(L, \mu)$.)

In order to prove the first of the two remaining claims, fix $0 \leq g \in C(L)$, and let $f \in C(K)$ and $u \in C(K)$ with $|u| \leq 1$. Then

$$\left| \int_L T f \cdot (gu) \, d\mu \right| = \left| \int_K f \cdot S(gu) \, d\nu \right| \leq \int_K |f| |S| g \, d\nu.$$

Taking the supremum over all these functions u , we obtain

$$\int_L |Tf| \cdot g \, d\mu \leq \int_K |f| |S|g \, d\nu.$$

This means that T extends to a contraction $T : L^1(K, (|S|g)\nu) \rightarrow L^1(L, g\mu)$. It follows that $|T|'_g(g\mu) \leq (|S|g)\nu$, where $|T|_g$ denotes the modulus of T considered as an operator $C(K) \rightarrow L^1(L, g\mu)$. However, since $L^1(L, \mu)$ “embeds” onto an ideal of $L^1(L, g\mu)$, it follows that $|T|_g = |T|$. Putting things together we obtain

$$\int_L |T|f \cdot g \, d\nu = \int_K f \, d|T|'_g(g\mu) \leq \int_K f \cdot |S|g \, d\nu$$

for $0 \leq f \in C(K)$. The converse inequality holds by symmetry, and the last remaining statement is obtained by integrating both measures against functions of the form $f \otimes g$. $\square$

Suppose that $T : C(K) \rightarrow L^1(L, \mu)$ is order-bounded. Then $|\mu_T| = \mu_{|T|}$ by Theorem 4.9, whence by standard integration theory there is a $\mu_{|T|}$ -almost everywhere unique $\lambda \in L^\infty(K \times L; \mu_{|T|})$ with $|\lambda| = 1$ almost everywhere and

$$\int_{K \times L} F(x, y) \, d\mu_T = \int_{K \times L} F(x, y) \lambda(x, y) \, d\mu_{|T|} \quad (4.5)$$

for all $F \in L^1(K \times L; \mu_{|T|})$. This leads to the following corollary for situation that $K = L$ and $\mu = \nu$.

Corollary 4.14. *Let (K, μ) be a compact measure space, let $T : C(K) \rightarrow L^1(K, \mu)$ be an order-bounded operator, and let $\lambda \in L^\infty(K \times K, \mu_{|T|})$ with $|\lambda| = 1$ almost everywhere and such that (4.5) holds for all $F \in L^1(K \times L; \mu_{|T|})$. Suppose, in addition, that T is symmetric, i.e., T satisfies*

$$\int_K Tf \cdot \bar{g} \, d\mu = \int_K f \cdot \overline{Tg} \, d\mu \quad (f, g \in C(K)).$$

Then $|T|$ is symmetric, too, and

$$\lambda(x, y) = \overline{\lambda(y, x)} \quad \text{for } \mu_{|T|}\text{-almost all } (x, y) \in K^2.$$

Proof. Note that, by hypothesis, (4.4) holds with $S = \overline{T}$, whence it holds for T and S replaced by $|T|$ and $|S| = |T|$, respectively. It follows that $|T|$ is symmetric and that $r_*\mu_{|T|} = \mu_{|T|}$. The last assertion is now straightforward. $\square$

The following is the main result of this section. It has essentially been proved by Carbonaro and Dragičević [2, p.22/23].

Theorem 4.15 (Disintegration). *Let (K, μ) be a compact measure space, and let T be a symmetric absolute contraction on $L^1(K, \mu)$. Then*

$$\int_K (I - T)f \cdot g \, d\mu = \int_K (I - M_{|T|1})f \cdot g \, d\mu$$

$$+ \int_{K \times K} \int_{Z_2} \left[\mathbf{I} - \begin{pmatrix} 0 & \overline{\lambda(x, y)} \\ \lambda(x, y) & 0 \end{pmatrix} \right] \begin{pmatrix} f(x) \\ f(y) \end{pmatrix} \cdot \begin{pmatrix} g(x) \\ g(y) \end{pmatrix} d\zeta_2 d\mu_{|T|}(x, y)$$

for all $f \in L^p(K, \mu)$, $g \in L^q(K, \mu)$, $1 \leq p \leq \infty$.

Proof. We first write $\mathbf{I} - T = (\mathbf{I} - M_{|T|\mathbf{1}}) + (M_{|T|\mathbf{1}} - T)$ and then compute

$$\begin{aligned} \int_K (M_{|T|\mathbf{1}} - T)f \cdot g d\mu &= \int_K (|T|\mathbf{1})f \cdot g d\mu - \int_K Tf \cdot g d\mu \\ &= \int_{K^2} \mathbf{1} \otimes fg d\mu_{|T|} - \int_{K^2} f \otimes g d\mu_T = \int_{K^2} \mathbf{1} \otimes fg - (f \otimes g)\bar{\lambda} d\mu_{|T|}. \end{aligned}$$

Since T is symmetric and $|\overline{T}| = |T|$, also $|T|$ is symmetric and $\mu_{|T|}$ is a symmetric positive measure. Therefore, by a change of variable $(x, y) \mapsto (y, x)$ in the formula from above,

$$\int_K (M_{|T|\mathbf{1}} - T)f \cdot g d\mu = \int_{K^2} fg \otimes \mathbf{1} - (g \otimes f)\bar{\lambda} d\mu_{|T|}.$$

Taking the arithmetic average of this and the previous form we obtain the claimed formula. $\square$

Corollary 4.16. *Let (K, μ) be a compact measure space, and let T be a symmetric sub-Markovian operator on $L^1(K, \mu)$. Then*

$$\begin{aligned} \int_K (\mathbf{I} - T)f \cdot g d\mu &= \int_K (\mathbf{1} - T\mathbf{1})f \cdot g d\mu \\ &\quad + \int_{K \times K} \int_{Z_2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} f(x) \\ f(y) \end{pmatrix} \cdot \begin{pmatrix} g(x) \\ g(y) \end{pmatrix} d\zeta_2 d\mu_T(x, y) \end{aligned}$$

for all $f \in L^p(K, \mu)$, $g \in L^q(K, \mu)$, $1 \leq p \leq \infty$.

5. Proof of the Main Results

Let us return to the proof of the main result, Theorem 2.2. By the reduction steps from Section 3, one can suppose from the start that $X = (K, \mu)$ is a compact measure space, $A = \mathbf{I} - T$ for some symmetric absolute contraction on $L^1(K, \mu)$. In particular, the Disintegration Theorem 4.15 is applicable.

Let, as in the hypothesis of Theorem 2.2, $1 \leq p < \infty$, $d, m \in \mathbb{N}$ and $F_j, G_j : K \rightarrow \mathbb{C}^d$ be measurable functions for $1 \leq j \leq m$. The assertion to prove is:

For all measurable functions $\mathbf{f} \in \mathcal{M}(K, \mu; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in L^p(K, \mu)$ and $G_j(\mathbf{f}) \in L^q(K, \mu)$ for all $1 \leq j \leq m$:

$$\sum_{j=1}^m \operatorname{Re} \int_K (\mathbf{I} - T)F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) d\mu \geq 0.$$

and we may suppose that this assertion holds when (K, μ) is replaced by Z_2 , and T is replaced by E_λ for each $\lambda \in \mathbb{T}$.

Lemma 5.1. *Under the given hypotheses,*

$$\operatorname{Re} \sum_{j=1}^m F_j(x) G_j(x) \geq 0 \quad \text{for all } x \in \mathbb{C}^d. \quad (5.1)$$

Proof. Note that the integral inequality is *convex* in T , and that it holds trivially for $T = \mathbf{I}$. Since it holds for each $T = E_\lambda$, $\lambda \in \mathbb{T}$, it also holds for $T = \frac{1}{2}E_1 + \frac{1}{2}E_{-1} = 0$. Given $(x_1, \dots, x_d) \in \mathbb{C}^d$, let $f_j := (x_j, x_j)^t \in \mathcal{M}(Z_2)$ and inserting this into the inequality with $T = 0$ on Z_2 yields the claim. $\square$

Suppose now that $\mathbf{f} \in \mathcal{M}(K, \mu; \mathbb{C}^d)$ such that $F_j(\mathbf{f}) \in L^p(K, \mu)$ and $G_j(\mathbf{f}) \in L^q(K, \mu)$. We can apply the Disintegration Theorem 4.15 and obtain, for each $j = 1, \dots, m$

$$\begin{aligned} \int_K (\mathbf{I} - T) F_j(\mathbf{f}) \cdot G_j(\mathbf{f}) \, d\mu &= \int_K (\mathbf{1} - |T| \mathbf{1}) F_j(\mathbf{f}) G_j(\mathbf{f}) \, d\mu \\ &+ \int_{K \times K} \int_{Z_2} (\mathbf{I} - E_{\lambda(x,y)}) \left(\frac{F_j(\mathbf{f}(x))}{F_j(\mathbf{f}(y))} \right) \cdot \left(\frac{G_j(\mathbf{f}(x))}{G_j(\mathbf{f}(y))} \right) \, d\zeta_2 \, d\mu_{|T|}(x, y). \end{aligned}$$

Now sum over j and take the real part. Finally, apply Lemma 5.1 for the first summand and the hypothesis over $E_{\lambda(x,y)}$ for the second to conclude that the result has to be ≥ 0 . Hence, Theorem 2.2 is completely proved.

The corresponding results for symmetric sub-Markovian and Markovian semigroups (Theorem 2.3, Theorem 2.4) are proved similarly. (Note that by the reduction steps in Section 3 one only needs to show the assertion for the case that $A = \mathbf{I} - T$ where T is a symmetric absolute contraction on a compact measure space (K, ν) , and T is sub-Markovian or Markovian, respectively.

In the sub-Markovian case (Theorem 2.3), the hypothesis tells in particular that the statement is true for $T = 0$ on Z_2 , hence (5.1) holds. Now apply Corollary 4.16 and proceed as before.

In the Markovian case, one has $T\mathbf{1} = \mathbf{1}$ and the first summand in the disintegration formula of Corollary 4.16 vanishes. This leads to Theorem 2.4. (Note that in the Markovian case, (5.1) is not a necessary condition any more.)

6. Application: The Sector of Analyticity

Let $(S_t)_{t \geq 0}$ be an absolute contraction semigroup over a measure space X , and let $1 < p < \infty$. As a consequence of the Lumer–Phillips theorem, the semigroup $(S_t)_{t \geq 0}$ extends to an analytic contraction semigroup on $L^p(X)$ defined on the sector

$$\Sigma_\varphi := \{z \in \mathbb{C} \setminus 0 : |\arg z| < \varphi\}$$

(where $0 < \varphi \leq \frac{\pi}{2}$) if and only if

$$\operatorname{Re} \int_X e^{\pm \varphi i} (Af) \cdot \overline{f} |f|^{p-2} \geq 0 \quad (6.1)$$

for all $f \in \text{dom}(A_p)$. For some time it had been an open question whether, in the case that $(S_t)_t$ is a *symmetric* contraction semigroup, inequality (6.1) must hold for the angle $\varphi = \varphi_p$, where

$$\varphi_p := \arccos \left| 1 - \frac{2}{p} \right| = \arctan \frac{2\sqrt{p-1}}{|p-2|} \quad (6.2)$$

for $1 < p < \infty$. Such a result had been first established by Bakry [1] for a certain subclass of sub-Markovian symmetric semigroups and later extended to all sub-Markovian symmetric semigroups by Liskevich and Perelmuter [17]. That proof was subsequently improved by Nagel and Voigt [19] and in that form became part of Chapter 3 in Ouhabaz' book [20]. The best general result for all symmetric contraction semigroups had for a long time been the one by Cowling [4], when Kriegler finally settled the case with a positive answer in [16]. Carbonaro and Dragičević showed in [2, Remark 35] that the optimal angle can be obtained also from their results.

We shall see in this section that the general symmetric case reduces to the same scalar inequality as the sub-Markovian case. We apply Theorem 2.2 with $d = m = 1$, $F(x) = x$ and $G(x) = e^{\pm i\varphi} \bar{x} |x|^{p-2}$ ($G(0) = 0$). This yields the inequality

$$\text{Re} \left(e^{\pm i\varphi} \begin{pmatrix} 1 & -\bar{\lambda} \\ -\lambda & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} \cdot_{Z_2} \begin{pmatrix} z |z|^{p-2} \\ w |w|^{p-2} \end{pmatrix} \right) \geq 0$$

for all choices of $z, w \in \mathbb{C}$ and $\lambda \in \mathbb{T}$. (Recall that $\cdot_{Z_2}$ denotes the sesquilinear inner product on $L^2(Z_2)$.) If we replace w by λw in this inequality, we obtain the equivalent inequality

$$\text{Re} \left(e^{\pm i\varphi} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix} \cdot_{Z_2} \begin{pmatrix} z |z|^{p-2} \\ w |w|^{p-2} \end{pmatrix} \right) \geq 0.$$

For $w = 0$ the inequality reduces to $|z|^p \cos \varphi \geq 0$, which poses no further restriction on φ . For $w \neq 0$ we can replace z by wz and find the equivalent inequality

$$\text{Re} \left(e^{\pm i\varphi} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} z \\ 1 \end{pmatrix} \cdot_{Z_2} \begin{pmatrix} z |z|^{p-2} \\ 1 \end{pmatrix} \right) \geq 0,$$

i.e.,

$$\text{Re} \left(e^{\pm i\varphi} (z - 1)(\bar{z} |z|^{p-2} - 1) \right) \geq 0.$$

Reformulating this as an inequality between real and imaginary part and letting $\varphi = \varphi_p$ as above reduces to the inequality (2.1) in [17] which is proven there. (Actually, our argument shows that the proof can be simplified since there is only one complex variable to deal with.)

Corollary 6.1 (Kriegler). *Let $-A$ be the generator of a symmetric contraction semigroup $S = (S_t)_{t \geq 0}$ over some measure space X , and let $1 < p < \infty$. Then S extends to an analytic semigroup of contractions on $L^p(X)$ on the sector Σ_{φ_p} .*

Appendix A. On Homomorphisms of Probability Spaces

Suppose that $X = (X, \Sigma, \mu)$ and $X' = (X', \Sigma', \mu')$ are probability spaces and

$$\Phi : L^1(X) \rightarrow L^1(X')$$

is a one-preserving isometric lattice homomorphism.² This means that Φ is an isometric embedding for the L^1 -norms, $\Phi(\mathbf{1}) = \mathbf{1}$ and $|\Phi f| = \Phi |f|$ for all $f \in L^1(X)$.

The positivity of Φ implies in particular that $\Phi(\overline{f}) = \overline{\Phi f}$ for all $f \in L^1(X)$. Finally,

$$\int_X f = \int_{X'} \Phi f$$

for all $f \in L^1(X)$, since this is true for all $f \geq 0$.

In this appendix we show how to (canonically) extend Φ to a homomorphic (as lattices and $*$ -algebras) embedding

$$\Phi : \mathcal{M}(X) \rightarrow \mathcal{M}(X')$$

where $\mathcal{M}(X)$ and $\mathcal{M}(X')$ denote the spaces of all measurable $\mathbb{C}$ -valued functions modulo almost everywhere equality on X and X' , respectively. Note that $\mathcal{M}(X)$ is a complete metric space with respect to the metric

$$d_X(f, g) := \int_X \frac{|f - g|}{1 + |f - g|}.$$

The following lemma is the key property.

Lemma A.1. *In the situation from above, Φ restricts to an embedding of C^* -algebras $\Phi : L^\infty(X) \rightarrow L^\infty(X')$. Moreover, for any $f \in L^1(X)$,*

$$\mu[|f| > 0] = \mu'[\Phi f > 0]$$

In particular, $[f = 0]$ is a μ -null set if and only if $[\Phi f = 0]$ is a μ' -null set.

Proof. It is clear that Φ restricts to a one-preserving isometric lattice homomorphism between the respective L^∞ -spaces. So only the multiplicativity $\Phi(fg) = (\Phi f)(\Phi g)$ is to be shown. This is well-known, see e.g. [5, Chap. 7], but we repeat the argument for the convenience of the reader. By bilinearity, it suffices to consider $f, g \geq 0$. Then, by polarization, it suffices to consider $f = g$, which reduces the problem to establish that $\Phi(f^2) = (\Phi f)^2$. Now, for any $x \geq 0$, $x^2 = \sup_{t \geq 0} 2tx - t^2$. Hence, $f^2 = \sup_{t \geq 0} 2tf - t^2 \mathbf{1}$ in the Banach lattice sense. But Φ is a lattice homomorphism and $\Phi \mathbf{1} = \mathbf{1}$, whence

$$\Phi(f^2) = \Phi\left(\sup_{t \geq 0} 2tf - t^2 \mathbf{1}\right) = \sup_{t \geq 0} 2t(\Phi f) - t^2 \mathbf{1} = (\Phi f)^2.$$

The remaining statement follows from:

$$\mu[|f| > 0] = \lim_{n \rightarrow \infty} \int_X (n|f| \wedge \mathbf{1}) = \lim_{n \rightarrow \infty} \int_{X'} \Phi(n|f| \wedge \mathbf{1})$$

²In [5, Chap. 12], this is called a *Markov embedding*. It is the functional-analytic analogue of a *factor map* (=homomorphism in the category of probability spaces) $X' \rightarrow X$.

$$= \lim_{n \rightarrow \infty} \int_{X'} n |\Phi f| \wedge \mathbf{1} = \mu' [|\Phi f| > 0]. \quad \square$$

Let $f \in \mathcal{M}(X)$. Then the function $e := \frac{1}{1+|f|}$ has the property that $e, ef \in L^\infty(X)$. Moreover, by Lemma A.1, $[\Phi e = 0]$ is μ' -null set. Hence, Φe is an invertible element in the algebra $\mathcal{M}(X')$, and we can define

$$\widehat{\Phi} f := \frac{\Phi(ef)}{\Phi e} \in \mathcal{M}(X').$$

Lemma A.2. *The so-defined mapping $\widehat{\Phi} : \mathcal{M}(X) \rightarrow \mathcal{M}(X')$ has the following properties:*

- a) $\widehat{\Phi}$ is an extension of Φ .
- b) $\widehat{\Phi}$ is a unital $*$ -algebra and lattice homomorphism.
- c) $\int_{X'} \widehat{\Phi} f = \int_X f$ whenever $0 \leq f \in \mathcal{M}(X)$.
- d) $\widehat{\Phi}$ is an isometry with respect to the canonical metrics d_X and $d_{X'}$.
- e) If Φ is bijective then so is $\widehat{\Phi}$.
- f) The mapping $\widehat{\Phi} : \mathcal{M}(X) \rightarrow \mathcal{M}(X')$ is uniquely determined by the property that it extends Φ and it is multiplicative, i.e., satisfies $\widehat{\Phi}(fg) = \widehat{\Phi} f \cdot \widehat{\Phi} g$ for all $f, g \in \mathcal{M}(X)$.

Proof. a) and b) This is straightforward and left to the reader.

c) By the monotone convergence theorem,

$$\begin{aligned} \int_X f &= \sup_{n \in \mathbb{N}} \int_X (f \wedge n\mathbf{1}) = \sup_{n \in \mathbb{N}} \int_X \Phi(f \wedge n\mathbf{1}) = \sup_{n \in \mathbb{N}} \int_{X'} \widehat{\Phi}(f \wedge n\mathbf{1}) \\ &= \sup_{n \in \mathbb{N}} \int_{X'} (\widehat{\Phi} f \wedge n\mathbf{1}) = \int_{X'} \widehat{\Phi} f. \end{aligned}$$

d) Follows from b) and c).

e) Suppose that $L^\infty(X') \subseteq \text{ran}(\Phi)$ and let $g \in \mathcal{M}(X')$ be arbitrary. Then, by Lemma A.1, there are $e, h \in L^\infty(X)$ such that

$$\Phi e = \frac{1}{1+|g|} \quad \text{and} \quad \Phi h = \frac{g}{1+|g|} = g \Phi e.$$

Again by Lemma A.1, $\mu[e = 0] = 0$, whence we can define $f := \frac{h}{e} \in \mathcal{M}(X)$. It follows that $\Phi f = g$.

f) Suppose that $\Psi : \mathcal{M}(X) \rightarrow \mathcal{M}(X')$ is multiplicative and extends Φ . Let $f \in \mathcal{M}(X)$ and define $e := \frac{1}{1+|f|}$ as before. Then $f, ef \in L^\infty(X)$ and hence

$$\Phi e \cdot \Psi f = \Psi e \cdot \Psi f = \Psi(ef) = \Phi(ef).$$

Since Φe is an invertible element in $\mathcal{M}(X')$ (as seen above), it follows that

$$\Psi f = \frac{\Phi(ef)}{\Phi e} = \widehat{\Phi} f$$

as claimed. $\square$

By abuse of notation, we write Φ again instead of $\widehat{\Phi}$. It is clear that Φ allows a further extension to $\mathbb{C}^d$ -valued functions by

$$\Phi(\mathbf{f}) = \Phi(f_1, \dots, f_d) := (\Phi f_1, \dots, \Phi f_d) \quad \text{for } \mathbf{f} = (f_1, \dots, f_d) \in \mathcal{M}(X; \mathbb{C}^d).$$

Now we are well-prepared for the final result of this appendix.

Theorem A.3. *Let X and X' be probability spaces, and let $\Phi : L^1(X) \rightarrow L^1(X')$ be a one-preserving isometric lattice isomorphism, with its canonical extension $\Phi : \mathcal{M}(X; \mathbb{C}^d) \rightarrow \mathcal{M}(X'; \mathbb{C}^d)$, $d \in \mathbb{N}$. Then*

$$\Phi(F(\mathbf{f})) = F(\Phi \mathbf{f}) \quad \text{almost everywhere} \quad (\text{A.1})$$

for every Borel measurable function $F : \mathbb{C}^d \rightarrow \mathbb{C}$ and every $\mathbf{f} \in \mathcal{M}(X; \mathbb{C}^d)$.

Proof. By linearity we may suppose that $F \geq 0$. Next, by approximating $F \wedge n\mathbf{1} \nearrow F$, we may suppose that F is bounded. Then F is a uniform limit of positive simple functions, whence we may suppose without loss of generality that $F = \mathbf{1}_B$, where B is a Borel set in $\mathbb{C}^d$. In this case, (A.1) becomes

$$\Phi(\mathbf{1}_{[(f_1, \dots, f_d) \in B]}) = \mathbf{1}_{[(\Phi f_1, \dots, \Phi f_d) \in B]} \quad \text{almost everywhere.}$$

Let $\mathcal{B}$ be the set of all Borel subsets of $\mathbb{C}^d$ that satisfy this. Then $\mathcal{B}$ is a Dynkin system, so it suffices to show that each rectangle is contained in $\mathcal{B}$. Since Φ is multiplicative, this reduces the case to $d = 1$, f is real valued and $B = (a, b]$. Now $[a < f \leq b] = [a < f] \cap [b < f]^c$, which reduces the situation to $B = (a, \infty)$. Now

$$\mathbf{1}_{[a < f]} = L^1\text{-}\lim_{n \rightarrow \infty} n(f - a\mathbf{1})^+ \wedge \mathbf{1},$$

and applying Φ concludes the proof. $\square$

Remarks A.4. 1) As a consequence of Theorem A.3, $\Phi|f|^p = |\Phi f|^p$ for any $f \in \mathcal{M}(X)$ and $p > 0$, so Φ restricts to an isometric isomorphism of L^p -spaces for each $p > 0$.

2) The extension of the original L^1 -isomorphism Φ to $\mathcal{M}(X)$ is uniquely determined by the requirement that Φ is continuous for the metrics d_X and $d_{X'}$.

3) One can extend Φ to a lattice homomorphism

$$\Phi : \mathcal{M}(X; [0, \infty]) \rightarrow \mathcal{M}(X'; [0, \infty])$$

by defining $\Phi f := \tau^{-1} \circ \Phi(\tau \circ f)$, where $\tau : [0, \infty] \rightarrow [0, 1]$ is any order-preserving bijection. Using this one can then show that Φ maps almost everywhere convergent sequences to almost everywhere convergent sequences.

Acknowledgement: I am very grateful to Tom ter Elst (Auckland) for his kind invitation in March/April 2014, and the pleasure of studying together the paper [2]. Furthermore, I thank Hendrik Vogt for some stimulating discussions and the anonymous referee for his careful reading of the manuscript and some very useful remarks.

References

- [1] BAKRY, D. Sur l'interpolation complexe des semigroupes de diffusion. In *Séminaire de Probabilités, XXIII*, vol. 1372 of *Lecture Notes in Math.* Springer, Berlin, 1989, pp. 1–20.
- [2] CARBONARO, A., AND DRAGICEVIC, O. Functional calculus for generators of symmetric contraction semigroups. arXiv:1308.1338v1.
- [3] CHACON, R. V., AND KRENGEL, U. Linear modulus of linear operator. *Proc. Amer. Math. Soc.* 15 (1964), 553–559.
- [4] COWLING, M. G. Harmonic analysis on semigroups. *Ann. of Math. (2)* 117, 2 (1983), 267–283.
- [5] EISNER, T., FARKAS, B., HAASE, M., AND NAGEL, R. *Operator Theoretic Aspects of Ergodic Theory*, vol. 272 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2015.
- [6] FUKUSHIMA, M., OSHIMA, Y., AND TAKEDA, M. *Dirichlet forms and symmetric Markov processes*, extended ed., vol. 19 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 2011.
- [7] GELFAND, I. On normed rings. *C. R. (Doklady) Acad. Sci. URSS, n. Ser.* 23 (1939), 430–432.
- [8] GELFAND, I., AND NEUMARK, M. On the imbedding of normed rings into the ring of operators in Hilbert space. *Rec. Math. [Mat. Sbornik] N.S.* 12(54) (1943), 197–213.
- [9] GROTHENDIECK, A. *Produits tensoriels topologiques et espaces nucléaires*, vol. 16 of *Mem. Amer. Math. Soc.* American Mathematical Society, Providence, 1955.
- [10] HAASE, M. Convexity inequalities for positive operators. *Positivity* 11, 1 (2007), 57–68.
- [11] HALMOS, P. R. On a theorem of Dieudonné. *Proc. Nat. Acad. Sci. U. S. A.* 35 (1949), 38–42.
- [12] HUANG, S.-Z. Inequalities for submarkovian operators and submarkovian semigroups. *Math. Nachr.* 243 (2002), 75–91.
- [13] KAKUTANI, S. Concrete representation of abstract (L) -spaces and the mean ergodic theorem. *Ann. of Math. (2)* 42 (1941), 523–537.
- [14] KAKUTANI, S. Concrete representation of abstract (M) -spaces. (A characterization of the space of continuous functions.). *Ann. of Math. (2)* 42 (1941), 994–1024.
- [15] KANTOROVITCH, L. Linear operations in semi-ordered spaces. I. *Rec. Math. Moscou, n. Ser.* 7 (1940), 209–284.
- [16] KRIEGLER, C. Analyticity angle for non-commutative diffusion semigroups. *J. Lond. Math. Soc. (2)* 83, 1 (2011), 168–186.
- [17] LISKEVICH, V. A., AND PEREL'MUTER, M. A. Analyticity of sub-Markovian semigroups. *Proc. Amer. Math. Soc.* 123, 4 (1995), 1097–1104.
- [18] LUXEMBURG, W. A. J., AND ZAAANEN, A. C. The linear modulus of an order bounded linear transformation. I & II. *Nederl. Akad. Wetensch. Proc. Ser. A* 74=Indag. Math. 33 (1971), 422–447.
- [19] NAGEL, R., AND VOIGT, J. On inequalities for symmetric sub-Markovian operators. *Arch. Math. (Basel)* 67, 4 (1996), 308–311.

- [20] OUHABAZ, E. M. *Analysis of heat equations on domains*, vol. 31 of *London Mathematical Society Monographs Series*. Princeton University Press, Princeton, NJ, 2005.
- [21] SCHAEFER, H. H. *Banach lattices and positive operators*. Springer-Verlag, New York-Heidelberg, 1974. Die Grundlehren der mathematischen Wissenschaften, Band 215.
- [22] SEGAL, I. E. Equivalences of measure spaces. *Amer. J. Math.* 73 (1951), 275–313.
- [23] STEIN, E. M. *Topics in harmonic analysis related to the Littlewood-Paley theory*. Annals of Mathematics Studies, No. 63. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970.
- [24] STONE, M. H. Applications of the theory of Boolean rings to general topology. *Trans. Amer. Math. Soc.* 41, 3 (1937), 375–481.
- [25] VOIGT, J. One-parameter semigroups acting simultaneously on different L_p -spaces. *Bull. Soc. Roy. Sci. Liège* 61, 6 (1992), 465–470.

Markus Haase
Christian-Albrechts Universität zu Kiel
Mathematisches Seminar
Ludewig-Meyn-Straße 4, 24098 Kiel, Germany
e-mail: haase@math.uni-kiel.de